

# Superlinear nonlocal fractional problems with infinitely many solutions

Zhang Binlin<sup>1</sup>, Giovanni Molica Bisci<sup>2</sup>  
and Raffaella Servadei<sup>3</sup>

<sup>1</sup> Department of Mathematics, Heilongjiang Institute of Technology, 150050 Harbin, People's Republic of China

<sup>2</sup> Dipartimento PAU, Università 'Mediterranea' di Reggio Calabria, Via Melissari 24, 89124 Reggio Calabria, Italy

<sup>3</sup> Dipartimento di Scienze di Base e Fondamenti, Università degli Studi di Urbino 'Carlo Bo', Piazza della Repubblica 13, 61029 Urbino (Pesaro e Urbino), Italy

E-mail: [zhbinlin@gmail.com](mailto:zhbinlin@gmail.com), [gmolica@unirc.it](mailto:gmolica@unirc.it) and [raffaella.servadei@uniurb.it](mailto:raffaella.servadei@uniurb.it)

Received 26 August 2014, revised 25 March 2015

Accepted for publication 22 April 2015

Published



CrossMark

Recommended by Dr Jean-Claude Saut

## Abstract

In this paper we study the existence of infinitely many weak solutions for equations driven by nonlocal integrodifferential operators with homogeneous Dirichlet boundary conditions. A model for these operators is given by the fractional Laplacian

$$-(-\Delta)^s u(x) := \int_{\mathbb{R}^n} \frac{u(x+y) + u(x-y) - 2u(x)}{|y|^{n+2s}} dy, \quad x \in \mathbb{R}^n$$

where  $s \in (0, 1)$  is fixed.

We consider different superlinear growth assumptions on the nonlinearity, starting from the well-known Ambrosetti–Rabinowitz condition. In this framework we obtain three different results about the existence of infinitely many weak solutions for the problem under consideration, by using the Fountain Theorem. All these theorems extend some classical results for semilinear Laplacian equations to the nonlocal fractional setting.

Keywords: fractional Laplacian, nonlocal problems, fountain theorem

Mathematics Subject Classification: 49J35, 35A15, 35S15, 47G20, 45G05

### 1. Introduction and main results

In this paper we are concerned with the existence of infinitely many weak solutions of the nonlocal fractional equations whose prototype of order  $s \in (0, 1)$  is given by

$$\begin{cases} (-\Delta)^s u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases} \tag{1.1}$$

which is the counterpart of this Laplace equation

$$\begin{cases} -\Delta u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{in } \partial\Omega. \end{cases} \tag{1.2}$$

Here  $(-\Delta)^s$  is the fractional Laplace operator, which, up to normalization factors, may be defined as

$$-(-\Delta)^s u(x) := \int_{\mathbb{R}^n} \frac{u(x+y) + u(x-y) - 2u(x)}{|y|^{n+2s}} dy, \quad x \in \mathbb{R}^n.$$

In recent years, a great attention has been focused on the study of fractional and nonlocal operators of elliptic type, both for the pure mathematical research and for concrete real-world applications. Fractional and nonlocal operators appear in many fields such as, among the others, optimization, finance, phase transitions, stratified materials, anomalous diffusion, crystal dislocation, soft thin films, semipermeable membranes, flame propagation, conservation laws, ultra-relativistic limits of quantum mechanics, quasi-geostrophic flows, multiple scattering, minimal surfaces, materials science and water waves. This is one of the reason why, recently, nonlocal fractional problems are widely studied in the literature in many different contexts. Just to name a few, we recall, for instance, the following papers and the references therein: [9,10,29] for regularity results, [4,5,8,17,24,25,36] for the existence of solutions, [26] for multiplicity of solutions and [18,27] for Kirchhoff nonlocal fractional problems.

In [33] the authors considered the following general nonlocal problem

$$\begin{cases} -\mathcal{L}_K u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega. \end{cases} \tag{1.3}$$

Here  $\Omega$  is an open bounded subset of  $\mathbb{R}^n$  with continuous boundary  $\partial\Omega$ ,  $n > 2s$ ,  $s \in (0, 1)$ , while  $\mathcal{L}_K$  is the integrodifferential operator defined as follows

$$\mathcal{L}_K u(x) := \int_{\mathbb{R}^n} (u(x+y) + u(x-y) - 2u(x)) K(y) dy, \quad x \in \mathbb{R}^n, \tag{1.4}$$

with the kernel  $K : \mathbb{R}^n \setminus \{0\} \rightarrow (0, +\infty)$  such that

$$mK \in L^1(\mathbb{R}^n), \text{ where } m(x) = \min\{|x|^2, 1\}; \tag{1.5}$$

$$\text{there exists } \theta > 0 \text{ such that } K(x) \geq \theta|x|^{-(n+2s)} \text{ for any } x \in \mathbb{R}^n \setminus \{0\}. \tag{1.6}$$

A prototype for  $K$  is given by the singular kernel  $K(x) = |x|^{-(n+2s)}$  which gives rise to the fractional Laplace operator  $(-\Delta)^s$ . In [33] an existence theorem for problem (1.3) has been proved by using the Mountain Pass Theorem, when the nonlinear term  $f$  has a superlinear and subcritical growth.

Motivated by an evident and increasing interest in the current literature on fractional elliptic problems, here we are interested in the existence of infinitely many weak solutions of problem (1.3) under the same superlinear growth assumptions on  $f$  adopted in [33], that is

$f : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  is a function verifying the following standard conditions

$$f \in C(\bar{\Omega} \times \mathbb{R}) \tag{1.7}$$

there exist  $a_1, a_2 > 0$  and  $q \in (2, 2^*)$ ,  $2^* = 2n/(n - 2s)$ , such that  $|f(x, t)| \leq a_1 + a_2|t|^{q-1}$  for any  $x \in \Omega, t \in \mathbb{R}$ ;  $\tag{1.8}$

there exist  $\mu > 2$  and  $r > 0$  such that for any  $x \in \Omega, t \in \mathbb{R}, |t| \geq r$   $\tag{1.9}$

$$0 < \mu F(x, t) \leq tf(x, t),$$

where the function  $F$  is the primitive of  $f$  with respect to the second variable, that is

$$F(x, t) = \int_0^t f(x, \tau) d\tau. \tag{1.10}$$

When looking for infinitely many solutions, it is natural requiring some symmetry on the nonlinearity. In the sequel we will assume that the following further assumption on  $f$  is satisfied

$$f(x, -t) = -f(x, t) \text{ for any } x \in \Omega, t \in \mathbb{R}. \tag{1.11}$$

As a model for  $f$  we can take the function  $f(x, t) = a(x)|t|^{q-2}t$ , with  $a \in C(\bar{\Omega})$  and  $q \in (2, 2^*)$ .

In the literature assumption (1.9) is well-known and it is called *Ambrosetti–Rabinowitz condition*, since it was originally introduced by Ambrosetti and Rabinowitz in [3], where, as an application of the famous Mountain Pass Theorem, they obtained the existence of nontrivial solutions of problem (1.2), under superlinear and subcritical growth conditions on the right-hand side.

A lot of works concerning superlinear elliptic boundary value problem have been written by using this usual Ambrosetti–Rabinowitz condition (see, for instance, [35,37] and the references therein), whose role consists in ensuring the boundedness of the Palais–Smale sequences of the energy functional associated with the problem under consideration.

The Ambrosetti–Rabinowitz condition is a superlinear growth assumption on the nonlinearity  $f$ . Indeed, from (1.9) it follows that for some  $a_3, a_4 > 0$

$$F(x, t) \geq a_3|t|^\mu - a_4 \text{ for any } (x, t) \in \bar{\Omega} \times \mathbb{R}, \tag{1.12}$$

see, for instance, [34, lemma 4]. However, there are many functions which are superlinear at infinity, but do not satisfy the Ambrosetti–Rabinowitz condition. At this purpose, we would note that from (1.12) and the fact that  $\mu > 2$ , it follows that

$$\lim_{|t| \rightarrow +\infty} \frac{F(x, t)}{|t|^2} = +\infty \text{ uniformly for any } x \in \bar{\Omega}. \tag{1.13}$$

Of course, also condition (1.13) characterizes the nonlinearity  $f$  to be superlinear at infinity. It is easily seen that the function

$$f(x, t) = t \log(1 + |t|) \tag{1.14}$$

verifies condition (1.13) and does not satisfy (1.12) (and so, as a consequence, does not verify (1.9)).

In order to study the superlinear problem (1.2) in [19] Jeanjean introduced the following assumption on  $f$ :

$$\begin{aligned} &\text{there exists } \gamma \geq 1 \text{ such that for any } x \in \Omega \\ &\mathcal{F}(x, t') \leq \gamma \mathcal{F}(x, t) \text{ for any } t, t' \in \mathbb{R} \text{ with } 0 < t' \leq t, \end{aligned} \tag{1.15}$$

where

$$\mathcal{F}(x, t) = \frac{1}{2} tf(x, t) - F(x, t). \tag{1.16}$$

It is easy to see that the function (1.14) satisfies also the condition (1.15).

In recent years, condition (1.15) was often applied to consider the existence of nontrivial solutions for the superlinear problem (1.2) without the Ambrosetti–Rabinowitz condition, for example, see [2, 15, 20–22]. For other papers treating superlinear problems without the Ambrosetti–Rabinowitz condition we refer to [13, 16, 19, 23, 30, 38, 39] and references therein.

According to [22, proposition 2.3], the condition (1.15) is weaker than the following assumption:

$$\text{the function } t \mapsto \frac{f(\cdot, t)}{t} \text{ is increasing in } t \geq 0 \text{ and decreasing in } t \leq 0. \tag{1.17}$$

However, both (1.15) and (1.17) are global conditions, and hence they are not very satisfactory. For this reason, we replace the condition (1.17) with the following local condition introduced by Liu in [21]:

$$\text{there exists } \bar{t} > 0 \text{ such that for any } x \in \Omega \tag{1.18}$$

$$\text{the function } t \mapsto \frac{f(x, t)}{t} \text{ is increasing in } t \geq \bar{t} \text{ and decreasing in } t \leq -\bar{t}.$$

After this overview on the assumptions on the nonlinearity  $f$ , we would note that problem (1.3) is variational in nature and the energy functional associated with it is given by the functional  $\mathcal{J}_K : X_0 \rightarrow \mathbb{R}$  defined as

$$\mathcal{J}_K(u) := \frac{1}{2} \int_{\mathbb{R}^n \times \mathbb{R}^n} |u(x) - u(y)|^2 K(x - y) dx dy - \int_{\Omega} F(x, u(x)) dx, \tag{1.19}$$

where  $F$  is the function defined in (1.10).

Now, we can state our main result.

**Theorem 1.** *Let  $s \in (0, 1)$ ,  $n > 2s$  and  $\Omega$  be an open bounded set of  $\mathbb{R}^n$  with continuous boundary. Let  $K : \mathbb{R}^n \setminus \{0\} \rightarrow (0, +\infty)$  be a function satisfying (1.5) and (1.6) and let  $f : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  be a function verifying (1.7), (1.8), (1.11) and*

- (a) (1.9) or
- (b) (1.13) and (1.15) or
- (c) (1.13) and (1.18).

*Then, the problem (1.3) has infinitely many solutions  $u_j \in X_0$ ,  $j \in \mathbb{N}$ , whose energy  $\mathcal{J}_K(u_j) \rightarrow +\infty$  as  $j \rightarrow +\infty$ .*

We would remark that, due to the symmetry assumption (1.11), if  $u$  is a weak solution of problem (1.3), then also  $-u$  does. Hence, our results assure the existence of infinitely many pairs  $\{u_j, -u_j\}_{j \in \mathbb{N}}$  of weak solutions for (1.3).

The strategy in order to get the multiplicity results stated here above consists in looking for infinitely many critical points for the Euler–Lagrange functional associated with problem (1.3), namely here we will apply the Fountain Theorem proved by Bartsch in [6]. As usual for critical point theorems, we have to study the compactness properties of the functional together with its geometric features. With respect to the compactness, we will prove that the functional satisfies the classical Palais–Smale condition when the nonlinearity verifies the Ambrosetti–Rabinowitz assumption, while, for a right-hand side satisfying other superlinear conditions (see (1.13) and (1.15) or (1.18)), the Cerami condition will be considered. In both cases the main difficulty relies in the proof of the boundedness of the Palais–Smale (or Cerami) sequence.

As for the geometry of the functional, we will show that it is negative in a ball of a suitable finite-dimensional subspace of  $X_0$  and positive in a ball of an infinite-dimensional subspace. For the negativity of the functional we will mainly use the equivalence of the norms in finite-dimensional spaces, while for the other geometric features we will need a more careful analysis, strictly related to the superlinear assumptions on the nonlinear term  $f$ .

Finally, we would note that theorem 1 represents the nonlocal counterpart of [6, theorem 3.7], [22, theorem 1.1] and [21, theorem 1.4]. We would also point out that in [25, theorem 3.1] the author proved the existence of infinitely many weak solutions of problem (1.3) requiring conditions (1.7)–(1.9) and (1.11), but exploiting a method different from the one used here and, precisely, a symmetric version of the Mountain Pass Theorem for even functionals.

In [34] the authors studied the nonlocal problem

$$\begin{cases} -\mathcal{L}_K u - \lambda u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases} \tag{1.20}$$

and proved the existence of a nontrivial solution for it, for any  $\lambda \in \mathbb{R}$ , as an application of the Mountain Pass Theorem and of the Linking Theorem (see [3, 28]).

With the same arguments used in the proof of theorem 1 and with some careful estimates of the term containing the parameter  $\lambda$ , it is easy to see that problem (1.20) admits infinitely many weak solutions, under all the different superlinear assumptions on  $f$  we considered above.

The paper is organized as follows. In section 2 we will present some necessary preliminary notions and results. In section 3 we will discuss the compactness properties of the energy functional associated with the problem under consideration. Finally, section 4 will be devoted to the proofs of the main result of the paper.

## 2. Preliminaries

In this section we give some preliminary results which will be used in the sequel.

### 2.1. The functional space $X_0$

Problems (1.1) and (1.3) have a variational nature and, in order to study them from this point of view, what we first need is to give a suitable variational formulation for them.

The natural spaces to look for solutions for these problems are the fractional Sobolev spaces. On the other hand, the variational formulation needs to encode the Dirichlet datum  $u = 0$  in  $\mathbb{R}^n \setminus \Omega$ . In order to overcome this problem, in [32] (see also [33, 34]) the authors considered a new functional space, denoted by  $X_0$ , which is inspired to the fractional Sobolev spaces (but it is not equivalent to them) and seems to be the good space for writing the variational formulation of our problems.

The space  $X_0$  is defined as

$$X_0 := \{g \in X : g = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\},$$

where the functional space  $X$  denotes the linear space of Lebesgue measurable functions from  $\mathbb{R}^n$  to  $\mathbb{R}$  such that the restriction to  $\Omega$  of any function  $g$  in  $X$  belongs to  $L^2(\Omega)$  and

the map  $(x, y) \mapsto (g(x) - g(y))\sqrt{K(x - y)}$  is in  $L^2((\mathbb{R}^n \times \mathbb{R}^n) \setminus (\mathcal{C}\Omega \times \mathcal{C}\Omega), dx dy)$

(here  $\mathcal{C}\Omega := \mathbb{R}^n \setminus \Omega$ ). Moreover,  $X_0$  is endowed with the norm

$$X_0 \ni g \mapsto \|g\|_{X_0} := \left( \int_{\mathbb{R}^n \times \mathbb{R}^n} |g(x) - g(y)|^2 K(x - y) dx dy \right)^{1/2} \tag{2.1}$$

and  $(X_0, \|\cdot\|_{X_0})$  is a Hilbert space (for this see [33, Lemma 7]), with scalar product

$$\langle u, v \rangle_{X_0} := \int_{\mathbb{R}^n \times \mathbb{R}^n} (u(x) - u(y))(v(x) - v(y)) K(x - y) \, dx \, dy. \tag{2.2}$$

The usual fractional Sobolev space  $H^s(\Omega)$  is endowed with the so-called *Gagliardo norm* (see, for instance [1, 14]) given by

$$\|g\|_{H^s(\Omega)} := \|g\|_{L^2(\Omega)} + \left( \int_{\Omega \times \Omega} \frac{|g(x) - g(y)|^2}{|x - y|^{n+2s}} \, dx \, dy \right)^{1/2}. \tag{2.3}$$

It is easy to see that, even in the model case in which  $K(x) = |x|^{-(n+2s)}$ , the norms in (2.1) and (2.3) are not the same: this makes the space  $X_0$  not equivalent to the usual fractional Sobolev spaces and the classical fractional Sobolev space approach not sufficient for studying our problem from a variational point of view.

Just for completeness, we would recall that both the spaces  $X$  and  $X_0$  are non-empty, since  $C_0^2(\Omega) \subseteq X_0$  (see [32, lemma 5.1]), and that for a general kernel  $K$  satisfying conditions (1.5) and (1.6), the following inclusion holds true

$$X_0 \subseteq \{g \in H^s(\mathbb{R}^n) : g = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\},$$

while, when  $K(x) = |x|^{-(n+2s)}$ , the space  $X_0$  can be characterized as follows

$$X_0 = \{g \in H^s(\mathbb{R}^n) : g = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\}.$$

For further details on  $X$  and  $X_0$  we refer to [32–34], where various properties of these spaces were proved. While, for more details on the fractional Sobolev spaces we refer to [14] and to the references therein.

In the sequel, we also need some properties of the spectrum of the operator  $-\mathcal{L}_K$  (for a complete study we refer to [31, proposition 2.3], and [34, proposition 9 and appendix A]). We recall that the eigenvalue problem

$$\begin{cases} -\mathcal{L}_K u = \lambda u & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega. \end{cases} \tag{2.4}$$

possesses a divergent sequence of positive eigenvalues

$$\lambda_1 < \lambda_2 \leq \dots \leq \lambda_k \leq \lambda_{k+1} \leq \dots,$$

whose corresponding eigenfunctions will be denoted by  $e_k$ . From [34, proposition 9], we know that  $\{e_k\}_{k \in \mathbb{N}}$  can be chosen in such a way that this sequence provides an orthonormal basis in  $L^2(\Omega)$  and an orthogonal basis in  $X_0$ .

### 2.2. Weak solutions and energy functional of the problem

Along this paper we are interested in the existence of infinitely many weak solutions for problem (1.3), that is on solutions of the following problem

$$\begin{cases} \int_{\mathbb{R}^n \times \mathbb{R}^n} (u(x) - u(y))(\varphi(x) - \varphi(y)) K(x - y) \, dx \, dy = \int_{\Omega} f(x, u(x)) \varphi(x) \, dx \\ u \in X_0, \end{cases} \quad \forall \varphi \in X_0 \tag{2.5}$$

which represents the Euler–Lagrange equation of the energy functional  $\mathcal{J}_K$  defined in (1.19). We would remark that  $\mathcal{J}_K \in C^1(X_0)$  thanks to the assumptions on  $f$  and also due to the embedding properties of  $X_0$  into the classical Lebesgue spaces (see [33, lemmas 6 and 8] and [34, lemma 9]).

In order to prove our main result, stated in theorem 1, we will apply the Fountain Theorem due to Bartsch (see [6]), which, under suitable compactness and geometric assumptions on a functional, provides the existence of an unbounded sequence of critical value for it.

### 3. Verification of the compactness conditions

The compactness assumption required by the Fountain Theorem is the well-known *Palais–Smale condition* (see, for instance, [35, 37] and references therein), which in our framework reads as follows:

$\mathcal{J}_K$  satisfies the *Palais–Smale compactness condition* at level  $c \in \mathbb{R}$   
 if any sequence  $\{u_j\}_{j \in \mathbb{N}}$  in  $X_0$  such that  
 $\mathcal{J}_K(u_j) \rightarrow c$  and  $\sup \left\{ \left| \langle \mathcal{J}'_K(u_j), \varphi \rangle \right| : \varphi \in X_0, \|\varphi\|_{X_0} = 1 \right\} \rightarrow 0$  as  $j \rightarrow +\infty$ ,  
 admits a subsequence strongly convergent in  $X_0$ .

In the case when the right-hand side in problem (1.3) satisfies conditions (1.8) and (1.9), in the sequel we will prove that the corresponding energy functional  $\mathcal{J}_K$  verifies the Palais–Smale condition. While, when removing the Ambrosetti–Rabinowitz condition (1.9) and replacing it with assumptions (1.13) and (1.15) or (1.18), we will show that  $\mathcal{J}_K$  verifies another compactness assumption, say the well-know *Cerami condition*, which in our setting can be written as follows:

$\mathcal{J}_K$  satisfies the *Cerami compactness condition* at level  $c \in \mathbb{R}$   
 if any sequence  $\{u_j\}_{j \in \mathbb{N}}$  in  $X_0$  such that  
 $\mathcal{J}_K(u_j) \rightarrow c$  and  $(1 + \|u_j\|) \sup \left\{ \left| \langle \mathcal{J}'_K(u_j), \varphi \rangle \right| : \varphi \in X_0, \|\varphi\|_{X_0} = 1 \right\} \rightarrow 0$   
 as  $j \rightarrow +\infty$ , admits a subsequence strongly convergent in  $X_0$ .

The Cerami condition was introduced by Cerami in [11, 12] as a weak version of the Palais–Smale condition. Hence, the Fountain Theorem holds true also under this compactness assumption. We would remark that if a functional satisfies the Palais–Smale condition or the Cerami condition, then it verifies the deformation condition, that is it fits with the requirements of the Deformation Theorem.

#### 3.1. Nonlinearities satisfying the Ambrosetti–Rabinowitz condition

In this framework we prove the following result about the compactness of the functional  $\mathcal{J}_K$ :

**Proposition 2.** *Let  $f : \overline{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  be a function verifying conditions (1.7)–(1.9). Then,  $\mathcal{J}_K$  satisfies the Palais–Smale condition at any level  $c \in \mathbb{R}$ .*

**Proof.** Let  $c \in \mathbb{R}$  and let  $\{u_j\}_{j \in \mathbb{N}}$  be a sequence in  $X_0$  such that

$$\mathcal{J}_K(u_j) \rightarrow c \tag{3.1}$$

and

$$\sup \left\{ \left| \langle \mathcal{J}'_K(u_j), \varphi \rangle \right| : \varphi \in X_0, \|\varphi\|_{X_0} = 1 \right\} \rightarrow 0 \tag{3.2}$$

as  $j \rightarrow +\infty$ .

We proceed by steps: first of all we show that the sequence  $\{u_j\}_{j \in \mathbb{N}}$  is bounded in  $X_0$  and then that it admits a subsequence strongly convergent in  $X_0$ .

**Step 1.** *The sequence  $\{u_j\}_{j \in \mathbb{N}}$  is bounded in  $X_0$ .* For any  $j \in \mathbb{N}$  by (3.1) and (3.2) it easily follows that there exists  $\kappa > 0$  such that

$$\left| \left\langle \mathcal{J}'_K(u_j), \frac{u_j}{\|u_j\|_{X_0}} \right\rangle \right| \leq \kappa$$

and

$$|\mathcal{J}_K(u_j)| \leq \kappa,$$

so that

$$\mathcal{J}_K(u_j) - \frac{1}{\mu} \langle \mathcal{J}'_K(u_j), u_j \rangle \leq \kappa (1 + \|u_j\|_{X_0}), \tag{3.3}$$

where  $\mu$  is the parameter given in (1.9).

By (1.8) and integrating it is easily seen that for any  $x \in \overline{\Omega}$  and for any  $t \in \mathbb{R}$

$$|F(x, t)| \leq a_1 |t| + \frac{a_2}{q} |t|^q. \tag{3.4}$$

Hence, by (3.4) and again (1.8) we have that for any  $j \in \mathbb{N}$

$$\begin{aligned} & \left| \int_{\Omega \cap \{|u_j| \leq r\}} \left( F(x, u_j(x)) - \frac{1}{\mu} f(x, u_j(x)) u_j(x) \right) dx \right| \\ & \leq \left( a_1 r + \frac{a_2}{q} r^q + \frac{a_1}{\mu} r + \frac{a_2}{\mu} r^q \right) |\Omega| =: \tilde{\kappa}. \end{aligned} \tag{3.5}$$

Thus, thanks to (1.9) and (3.5), we get that

$$\begin{aligned} \mathcal{J}_K(u_j) - \frac{1}{\mu} \langle \mathcal{J}'_K(u_j), u_j \rangle &= \left( \frac{1}{2} - \frac{1}{\mu} \right) \|u_j\|_{X_0}^2 \\ &\quad - \int_{\Omega} \left( F(x, u_j(x)) - \frac{1}{\mu} f(x, u_j(x)) u_j(x) \right) dx \\ &\geq \left( \frac{1}{2} - \frac{1}{\mu} \right) \|u_j\|_{X_0}^2 \\ &\quad - \int_{\Omega \cap \{|u_j| \leq r\}} \left( F(x, u_j(x)) - \frac{1}{\mu} f(x, u_j(x)) u_j(x) \right) dx \\ &\geq \left( \frac{1}{2} - \frac{1}{\mu} \right) \|u_j\|_{X_0}^2 - \tilde{\kappa} \end{aligned} \tag{3.6}$$

for any  $j \in \mathbb{N}$ .

By (3.3), (3.6) and the fact that  $\mu > 2$  we have that

$$\left( \frac{1}{2} - \frac{1}{\mu} \right) \|u_j\|_{X_0}^2 \leq \kappa (1 + \|u_j\|_{X_0}) + \tilde{\kappa}$$

for any  $j \in \mathbb{N}$ . Hence, step 1 is proved.

**Step 2.** Up to a subsequence,  $\{u_j\}_{j \in \mathbb{N}}$  strongly converges in  $X_0$ . Since  $\{u_j\}_{j \in \mathbb{N}}$  is bounded in  $X_0$  by step 1 and  $X_0$  is a reflexive space (being a Hilbert space, by [33, lemma 7]), up to a subsequence, still denoted by  $\{u_j\}_{j \in \mathbb{N}}$ , there exists  $u_\infty \in X_0$  such that

$$\int_{\mathbb{R}^n \times \mathbb{R}^n} (u_j(x) - u_j(y))(\varphi(x) - \varphi(y))K(x - y) dx dy \rightarrow \tag{3.7}$$

$$\int_{\mathbb{R}^n \times \mathbb{R}^n} (u_\infty(x) - u_\infty(y))(\varphi(x) - \varphi(y))K(x - y) dx dy \quad \text{for any } \varphi \in X_0$$

as  $j \rightarrow +\infty$ . Moreover, up to a subsequence,

$$\begin{aligned} u_j &\rightarrow u_\infty \quad \text{in } L^q(\mathbb{R}^n) \\ u_j &\rightarrow u_\infty \quad \text{a.e. in } \mathbb{R}^n \end{aligned} \tag{3.8}$$

as  $j \rightarrow +\infty$  and there exists  $\ell \in L^q(\mathbb{R}^n)$  such that

$$|u_j(x)| \leq \ell(x) \quad \text{a.e. in } \mathbb{R}^n \quad \text{for any } j \in \mathbb{N} \tag{3.9}$$

(see, for instance, [7, theorem 4.9]).



By (1.8), (3.7)–(3.9), the fact that the map  $t \mapsto f(\cdot, t)$  is continuous in  $t \in \mathbb{R}$  and the Dominated Convergence Theorem we get

$$\int_{\Omega} f(x, u_j(x))u_j(x) \, dx \rightarrow \int_{\Omega} f(x, u_{\infty}(x))u_{\infty}(x) \, dx \tag{3.10}$$

and

$$\int_{\Omega} f(x, u_j(x))u_{\infty}(x) \, dx \rightarrow \int_{\Omega} f(x, u_{\infty}(x))u_{\infty}(x) \, dx \tag{3.11}$$

as  $j \rightarrow +\infty$ . Moreover, by (3.2) and step 1 we have that

$$0 \leftarrow \langle \mathcal{J}'_K(u_j), u_j \rangle = \int_{\mathbb{R}^n \times \mathbb{R}^n} |u_j(x) - u_j(y)|^2 K(x - y) \, dx \, dy - \int_{\Omega} f(x, u_j(x))u_j(x) \, dx$$

so that, by (3.10) we deduce that

$$\int_{\mathbb{R}^n \times \mathbb{R}^n} |u_j(x) - u_j(y)|^2 K(x - y) \, dx \, dy \rightarrow \int_{\Omega} f(x, u_{\infty}(x))u_{\infty}(x) \, dx \tag{3.12}$$

as  $j \rightarrow +\infty$ . Furthermore, again by (3.2), we get

$$\begin{aligned} 0 \leftarrow \langle \mathcal{J}'_K(u_j), u_{\infty} \rangle &= \int_{\mathbb{R}^n \times \mathbb{R}^n} (u_j(x) - u_j(y))(u_{\infty}(x) - u_{\infty}(y))K(x - y) \, dx \, dy \\ &\quad - \int_{\Omega} f(x, u_j(x))u_{\infty}(x) \, dx \end{aligned} \tag{3.13}$$

as  $j \rightarrow +\infty$ . By (3.7) with  $\varphi = u_{\infty}$ , (3.11) and (3.13) we obtain

$$\int_{\mathbb{R}^n \times \mathbb{R}^n} |u_{\infty}(x) - u_{\infty}(y)|^2 K(x - y) \, dx \, dy = \int_{\Omega} f(x, u_{\infty}(x))u_{\infty}(x) \, dx. \tag{3.14}$$

Thus, (3.12) and (3.14) give that

$$\|u_j\|_{X_0} \rightarrow \|u_{\infty}\|_{X_0}, \tag{3.15}$$

as  $j \rightarrow \infty$ .

Finally, it is easy to see that

$$\begin{aligned} \|u_j - u_{\infty}\|_{X_0}^2 &= \|u_j\|_{X_0}^2 + \|u_{\infty}\|_{X_0}^2 - 2 \int_{\mathbb{R}^n \times \mathbb{R}^n} (u_j(x) - u_j(y))(u_{\infty}(x) - u_{\infty}(y))K(x - y) \, dx \, dy \\ &\rightarrow 2\|u_{\infty}\|_{X_0}^2 - 2 \int_{\mathbb{R}^n \times \mathbb{R}^n} |u_{\infty}(x) - u_{\infty}(y)|^2 K(x - y) \, dx \, dy = 0 \end{aligned}$$

as  $j \rightarrow +\infty$ , thanks to (3.7) and (3.15). Then, the assertion of step 2 is proved. This concludes the proof of proposition 2.  $\square$

### 3.2. Nonlinearities under the superlinear assumptions (1.13) and (1.15)

In this framework we show that the functional  $\mathcal{J}_K$  verifies the Cerami condition. Before proving this fact, we would note that, as a consequence of the assumptions (1.11) and (1.15), the following condition is verified:

$$\begin{aligned} &\text{there exists } \gamma \geq 1 \text{ such that for any } x \in \Omega \\ &\mathcal{F}(x, t') \leq \gamma \mathcal{F}(x, t) \text{ for any } t, t' \in \mathbb{R} \text{ with } 0 < |t'| \leq |t|, \end{aligned} \tag{3.16}$$

where  $\mathcal{F}$  is the function given in (1.16).

Now, we are ready to prove the next result, that is

**Proposition 3.** *Let  $f : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  be a function verifying conditions (1.7), (1.8), (1.11), (1.13) and (1.15). Then,  $\mathcal{J}_K$  satisfies the Cerami condition at any level  $c \in \mathbb{R}$ .*

**Proof.** Let  $c \in \mathbb{R}$  and let  $\{u_j\}_{j \in \mathbb{N}}$  be a Cerami sequence in  $X_0$ , that is let  $\{u_j\}_{j \in \mathbb{N}}$  be such that

$$\mathcal{J}_K(u_j) \rightarrow c \tag{3.17}$$

and

$$(1 + \|u_j\|) \sup \left\{ |\langle \mathcal{J}'_K(u_j), \varphi \rangle| : \varphi \in X_0, \|\varphi\|_{X_0} = 1 \right\} \rightarrow 0 \tag{3.18}$$

as  $j \rightarrow +\infty$ .

First of all, we show that the sequence  $\{u_j\}_{j \in \mathbb{N}}$  is bounded in  $X_0$ . To this purpose we argue as in the proof of [15, lemma 2.2]. We assume, by contradiction, that  $\{u_j\}_{j \in \mathbb{N}}$  is unbounded in  $X_0$ , that is we may suppose that, up to a subsequence (still denoted by  $\{u_j\}_{j \in \mathbb{N}}$ )

$$\|u_j\|_{X_0} \rightarrow +\infty \tag{3.19}$$

as  $j \rightarrow +\infty$ .

Note that, by (3.18) and (3.19), it is easily seen that

$$\sup \left\{ |\langle \mathcal{J}'_K(u_j), \varphi \rangle| : \varphi \in X_0, \|\varphi\|_{X_0} = 1 \right\} \rightarrow 0 \tag{3.20}$$

as  $j \rightarrow +\infty$ .

Now, for any  $j \in \mathbb{N}$ , let

$$v_j = \frac{u_j}{\|u_j\|_{X_0}}. \tag{3.21}$$

Of course, the sequence  $\{v_j\}_{j \in \mathbb{N}}$  is bounded in  $X_0$  and so, up to a subsequence, we have that there exists  $v_\infty \in X_0$  such that

$$\begin{aligned} v_j &\rightarrow v_\infty \quad \text{in } L^q(\mathbb{R}^n) \\ v_j &\rightarrow v_\infty \quad \text{a.e. in } \mathbb{R}^n \end{aligned} \tag{3.22}$$

as  $j \rightarrow +\infty$  and there exists  $\ell \in L^q(\mathbb{R}^n)$  such that

$$|v_j(x)| \leq \ell(x) \quad \text{a.e. in } \mathbb{R}^n \quad \text{for any } j \in \mathbb{N} \tag{3.23}$$

(see [7, theorem 4.9]). In the sequel we will consider separately the cases when  $v_\infty \equiv 0$  and  $v_\infty \not\equiv 0$  and we will prove that in both these situations a contradiction occurs.

Firstly, let us suppose that

$$v_\infty \equiv 0. \tag{3.24}$$

As in [19], we can say that for any  $j \in \mathbb{N}$  there exists  $t_j \in [0, 1]$  such that

$$\mathcal{J}_K(t_j u_j) = \max_{t \in [0,1]} \mathcal{J}_K(t u_j). \tag{3.25}$$

Since (3.19) holds true, for any  $m \in \mathbb{N}$ , we can choose  $r_m = 2\sqrt{m}$  such that

$$r_m \|u_j\|_{X_0}^{-1} \in (0, 1), \tag{3.26}$$

provided  $j$  is large enough, say  $j > \bar{j}$ , with  $\bar{j} = \bar{j}(m)$ .

By (3.22) and the continuity of the function  $F$ , we get that

$$F(x, r_m v_j(x)) \rightarrow F(x, r_m v_\infty(x)) \quad \text{a.e. } x \in \Omega \tag{3.27}$$

as  $j \rightarrow +\infty$  for any  $m \in \mathbb{N}$ . Moreover, integrating (1.8) and taking into account (3.23), we have that

$$\begin{aligned} |F(x, r_m v_j(x))| &\leq a_1 |r_m v_j(x)| + \frac{a_2}{q} |r_m v_j(x)|^q \\ &\leq a_1 r_m \ell(x) + \frac{a_2}{q} (r_m \ell(x))^q \in L^1(\Omega), \end{aligned} \tag{3.28}$$

a.e.  $x \in \Omega$  and for any  $m, j \in \mathbb{N}$ . Hence, (3.27), (3.28) and the Dominated Convergence Theorem yield that

$$F(\cdot, r_m v_j(\cdot)) \rightarrow F(\cdot, r_m v_\infty(\cdot)) \quad \text{in } L^1(\Omega) \tag{3.29}$$

as  $j \rightarrow +\infty$  for any  $m \in \mathbb{N}$ . Since  $F(x, 0) = 0$  for any  $x \in \bar{\Omega}$  and (3.24) holds true, (3.29) gives that

$$\int_{\Omega} F(x, r_m v_j(x)) \, dx \rightarrow 0 \tag{3.30}$$

as  $j \rightarrow +\infty$  for any  $m \in \mathbb{N}$ . Thus, (3.25), (3.26) and (3.30) yield

$$\begin{aligned} \mathcal{J}_K(t_j u_j) &\geq \mathcal{J}_K(r_m \|u_j\|_{X_0}^{-1} u_j) \\ &= \mathcal{J}_K(r_m v_j) \\ &= \frac{1}{2} \|r_m v_j\|_{X_0}^2 - \int_{\Omega} F(x, r_m v_j(x)) \, dx \\ &= 2m - \int_{\Omega} F(x, r_m v_j(x)) \, dx \geq m, \end{aligned}$$

provided  $j$  is large enough and for any  $m \in \mathbb{N}$ . From this we deduce that

$$\mathcal{J}_K(t_j u_j) \rightarrow +\infty \tag{3.31}$$

as  $j \rightarrow +\infty$ .

Now, note that  $\mathcal{J}_K(0) = 0$  and (3.17) holds true. Combining these two facts it is easily seen that  $t_j \in (0, 1)$  and so, by (3.25), we get that

$$\left. \frac{d}{dt} \right|_{t=t_j} \mathcal{J}_K(t u_j) = 0$$

for any  $j \in \mathbb{N}$ . As a consequence of this, we have that

$$\langle \mathcal{J}'_K(t_j u_j), t_j u_j \rangle = t_j \left. \frac{d}{dt} \right|_{t=t_j} \mathcal{J}_K(t u_j) = 0. \tag{3.32}$$

We claim that

$$\limsup_{j \rightarrow +\infty} \mathcal{J}_K(t_j u_j) \leq \kappa, \tag{3.33}$$

for a suitable positive constant  $\kappa$ . Indeed, by (3.32) and using (3.16), we get

$$\begin{aligned} \frac{1}{\gamma} \mathcal{J}_K(t_j u_j) &= \frac{1}{\gamma} \left( \mathcal{J}_K(t_j u_j) - \frac{1}{2} \langle \mathcal{J}'_K(t_j u_j), t_j u_j \rangle \right) \\ &= \frac{1}{\gamma} \left( - \int_{\Omega} F(x, t_j u_j(x)) \, dx + \frac{1}{2} \int_{\Omega} t_j u_j(x) f(x, t_j u_j(x)) \, dx \right) \\ &= \frac{1}{\gamma} \int_{\Omega} \mathcal{F}(x, t_j u_j(x)) \, dx \\ &\leq \int_{\Omega} \mathcal{F}(x, u_j(x)) \, dx \end{aligned}$$

$$\begin{aligned} &= \int_{\Omega} \left[ \frac{1}{2} u_j(x) f(x, u_j(x)) - F(x, u_j(x)) \right] dx \\ &= \mathcal{J}_K(u_j) - \frac{1}{2} \langle \mathcal{J}'_K(u_j), u_j \rangle \rightarrow c \end{aligned}$$

as  $j \rightarrow +\infty$ , thanks to (3.17) and (3.20). This proves (3.33), which contradicts (3.31). Thus, the sequence  $\{u_j\}_{j \in \mathbb{N}}$  has to be bounded in  $X_0$ .

Now, suppose that  $v_{\infty} \neq 0$ . Then, the set  $\Omega' := \{x \in \Omega : v_{\infty}(x) \neq 0\}$  has positive Lebesgue measure and

$$|u_j(x)| \rightarrow +\infty \text{ a.e. } x \in \Omega' \tag{3.34}$$

as  $j \rightarrow +\infty$ , thanks to (3.21), (3.22) and the fact that  $v_{\infty} \neq 0$ .

By (3.17) and (3.19) it is easy to see that

$$\frac{\mathcal{J}_K(u_j)}{\|u_j\|_{X_0}^2} \rightarrow 0,$$

that is

$$\frac{1}{2} - \int_{\Omega'} \frac{F(x, u_j(x))}{\|u_j\|_{X_0}^2} dx - \int_{\Omega \setminus \Omega'} \frac{F(x, u_j(x))}{\|u_j\|_{X_0}^2} dx = o(1) \tag{3.35}$$

as  $j \rightarrow +\infty$ .

Let us consider separately the two integrals in formula (3.35). With respect to the first one, we have that

$$\begin{aligned} \frac{F(x, u_j(x))}{\|u_j\|_{X_0}^2} &= \frac{F(x, u_j(x)) |u_j(x)|^2}{|u_j(x)|^2 \|u_j\|_{X_0}^2} \\ &= \frac{F(x, u_j(x))}{|u_j(x)|^2} |v_j(x)|^2 \rightarrow +\infty \text{ a.e. } x \in \Omega' \end{aligned}$$

as  $j \rightarrow +\infty$ , thanks to (1.13), (3.22), (3.34) and the definition of  $\Omega'$ . Hence, by using the Fatou lemma, we obtain

$$\int_{\Omega'} \frac{F(x, u_j(x))}{\|u_j\|_{X_0}^2} dx \rightarrow +\infty \tag{3.36}$$

as  $j \rightarrow +\infty$ .

As for the second integral in (3.35), we claim that

$$\int_{\Omega \setminus \Omega'} \frac{F(x, u_j(x))}{\|u_j\|_{X_0}^2} dx \geq -\frac{\kappa}{\|u_j\|_{X_0}^2} |\Omega \setminus \Omega'|, \tag{3.37}$$

for some positive constant  $\kappa$ . Indeed by (1.13), it follows that

$$\lim_{|t| \rightarrow +\infty} F(x, t) = +\infty \text{ uniformly for any } x \in \overline{\Omega}. \tag{3.38}$$

Hence, by (3.38) there exist two positive constants  $\tilde{t}$  and  $H$  such that

$$F(x, t) \geq H \tag{3.39}$$

for every  $x \in \overline{\Omega}$  and  $|t| > \tilde{t}$ . On the other hand, since  $F$  is continuous in  $\overline{\Omega} \times \mathbb{R}$ , one has

$$F(x, t) \geq \min_{(x,t) \in \overline{\Omega} \times [-\tilde{t}, \tilde{t}]} F(x, t), \tag{3.40}$$

for every  $x \in \bar{\Omega}$  and  $|t| \leq \bar{t}$ . Note that  $\min_{(x,t) \in \bar{\Omega} \times [-\bar{t}, \bar{t}]} F(x, t) \leq 0$ , being  $F(x, 0) = 0$  for any  $x \in \bar{\Omega}$ . Then, by (3.39) and (3.40) it follows that

$$F(x, t) \geq -\kappa \quad \text{for any } (x, t) \in \bar{\Omega} \times \mathbb{R} \tag{3.41}$$

for some positive constant  $\kappa$ . Inequality (3.41) immediately yields the claim (3.37).

As a consequence of (3.19) and (3.37) it is easy to see that

$$\lim_{j \rightarrow +\infty} \int_{\Omega \setminus \Omega'} \frac{F(x, u_j(x))}{\|u_j\|_{X_0}^2} dx \geq 0, \tag{3.42}$$

(note that this limit exists thanks to (3.35) and (3.36)). All in all, by (3.35), (3.36) and (3.42) we get a contradiction. Thus, the sequence  $\{u_j\}_{j \in \mathbb{N}}$  is bounded in  $X_0$ .

In order to prove the assertion of proposition 3 from now on we can argue as in step 2 of the proof of proposition 2.  $\square$

We would remark that along the proof of proposition 3 the assumption (1.15) was used (and was crucial) just for proving the inequality (3.33).

### 3.3. Nonlinearities verifying the superlinear conditions (1.13) and (1.18)

In this setting we need the following lemma, which will be crucial in the proof of the main result of this subsection.

**Lemma 4.** *If (1.18) holds true, then for any  $x \in \Omega$ , the function  $\mathcal{F}(x, t)$  is increasing in  $t \geq \bar{t}$  and decreasing in  $t \leq -\bar{t}$ , where  $\mathcal{F}$  is the function given in (1.16).*

*In particular, there exists  $C_1 > 0$  such that*

$$\mathcal{F}(x, s) \leq \mathcal{F}(x, t) + C_1$$

*for any  $x \in \Omega$  and  $0 \leq s \leq t$  or  $t \leq s \leq 0$ .*

See [21, lemma 2.3] for details.

**Proposition 5.** *Let  $f : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  be a function verifying conditions (1.7), (1.8), (1.13) and (1.18). Then,  $\mathcal{J}_K$  satisfies the Cerami condition at any level  $c \in \mathbb{R}$ .*

**Proof.** We can argue exactly as in the proof of proposition 3. We just have to modify the proof of inequality (3.33): indeed, for proving it, in proposition 3 we used condition (1.15) (actually (3.16)), which now is no more assumed.

Here we will show the validity of (3.33) making use of the assumption (1.18) and of lemma 4. We point out that our notation is the one used in the proof of proposition 3. In view of Lemma 4 we have that

$$\begin{aligned} \mathcal{J}_K(t_j u_j) &= \mathcal{J}_K(t_j u_j) - \frac{1}{2} \langle \mathcal{J}'_K(t_j u_j), t_j u_j \rangle \\ &= \int_{\Omega} \mathcal{F}(x, t_j u_j(x)) dx \\ &= \int_{\{u_j \geq 0\}} \mathcal{F}(x, t_j u_j(x)) dx + \int_{\{u_j < 0\}} \mathcal{F}(x, t_j u_j(x)) dx \\ &\leq \int_{\{u_j \geq 0\}} [\mathcal{F}(x, u_j(x)) + C_1] + \int_{\{u_j < 0\}} [\mathcal{F}(x, u_j(x)) + C_1] \\ &= \int_{\Omega} \mathcal{F}(x, u_j(x)) dx + C_1 |\Omega| \\ &= \mathcal{J}_K(u_j) - \frac{1}{2} \langle \mathcal{J}'_K(u_j), u_j \rangle + C_1 |\Omega| \rightarrow c + C_1 |\Omega| \end{aligned}$$

as  $j \rightarrow +\infty$ . This proves (3.33). The proof of proposition 5 is thus completed.  $\square$

#### 4. The proof of the main result

In this section we give the proofs of the existence of infinitely many solutions for problem (1.3), both when the right-hand side satisfies the Ambrosetti–Rabinowitz condition and when other superlinear assumptions are required. In both cases the strategy consists in applying the Fountain Theorem of Bartsch (see [6]) to the functional  $\mathcal{J}_K$ .

Following the notation used in [6, theorem 2.5] (see also [37]), in the sequel for any  $k \in \mathbb{N}$  we put

$$Y_k := \text{span}\{e_1, \dots, e_k\}$$

and

$$Z_k := \overline{\text{span}\{e_k, e_{k+1}, \dots\}}.$$

Since  $Y_k$  is finite-dimensional, all norms on  $Y_k$  are equivalent. Therefore, there exist two positive constants  $C_{k,q}$  and  $\tilde{C}_{k,q}$ , depending on  $k$  and  $q$ , such that for any  $u \in Y_k$

$$C_{k,q} \|u\|_{X_0} \leq \|u\|_{L^q(\Omega)} \leq \tilde{C}_{k,q} \|u\|_{X_0}. \tag{4.1}$$

The Fountain Theorem provides the existence of an unbounded sequence of critical value for a smooth functional, under suitable compactness condition (say, the Palais–Smale condition) and geometric assumptions on it, which, in our framework, read as follows:

- (i)  $a_k := \max \left\{ \mathcal{J}_K(u) : u \in Y_k, \|u\|_{X_0} = r_k \right\} \leq 0$ ;
- (ii)  $b_k := \inf \left\{ \mathcal{J}_K(u) : u \in Z_k, \|u\|_{X_0} = \gamma_k \right\} \rightarrow \infty$  as  $k \rightarrow \infty$ .

##### 4.1. Proof of theorem 1 under assumption (a)

In order to perform the proof of theorem 1 when condition (a) is assumed, we first need the following result:

**Lemma 6.** *Let  $1 \leq q < 2^*$  and, for any  $k \in \mathbb{N}$ , let*

$$\beta_k := \sup \left\{ \|u\|_{L^q(\Omega)} : u \in Z_k, \|u\|_{X_0} = 1 \right\}.$$

*Then,  $\beta_k \rightarrow 0$  as  $k \rightarrow \infty$ .*

**Proof.** By definition of  $Z_k$ , we have that  $Z_{k+1} \subset Z_k$  and so, as a consequence,  $0 < \beta_{k+1} \leq \beta_k$  for any  $k \in \mathbb{N}$ . Hence,

$$\beta_k \rightarrow \beta \tag{4.2}$$

as  $k \rightarrow +\infty$ , for some  $\beta \geq 0$ . Moreover, by definition of  $\beta_k$ , for any  $k \in \mathbb{N}$  there exists  $u_k \in Z_k$  such that

$$\|u_k\|_{X_0} = 1 \quad \text{and} \quad \|u_k\|_{L^q(\Omega)} > \beta_k/2. \tag{4.3}$$

Since  $X_0$  is a Hilbert space, and hence a reflexive Banach space, there exist  $u_\infty \in X_0$  and a subsequence of  $u_k$  (still denoted by  $u_k$ ) such that  $u_k \rightarrow u_\infty$  weakly in  $X_0$ , that is

$$\langle u_k, \varphi \rangle_{X_0} \rightarrow \langle u_\infty, \varphi \rangle_{X_0} \quad \text{for any } \varphi \in X_0$$

as  $k \rightarrow +\infty$ . Since each  $Z_k$  is convex and closed, hence it is closed for the weak topology. Consequently,

$$u_\infty \in \bigcap_{k=1}^{+\infty} Z_k = \{0\}.$$

Hence, by the Sobolev embedding theorem, we get

$$u_k \rightarrow 0 \quad \text{in } L^q(\Omega) \tag{4.4}$$

as  $k \rightarrow +\infty$ . By (4.2), the fact that  $\beta$  is nonnegative, (4.3) and (4.4) we get that  $\beta_k \rightarrow 0$  as  $k \rightarrow +\infty$  and this concludes the proof of lemma 6.  $\square$

**Proof of theorem 1 under assumption (a).** We mimic the proof of [37, theorem 3.7]. By proposition 2 we have that  $\mathcal{J}_K$  satisfies the Palais–Smale condition, while, by (1.11) we get that  $\mathcal{J}_K(-u) = \mathcal{J}_K(u)$  for any  $u \in X_0$ . In order to apply the Fountain Theorem, it remains to study the geometry of the functional  $\mathcal{J}_K$ . To this purpose, let us proceed by steps.

**Step 1.** For any  $k \in \mathbb{N}$  there exists  $r_k > 0$  such that

$$a_k = \max \left\{ \mathcal{J}_K(u) : u \in Y_k, \|u\|_{X_0} = r_k \right\} \leq 0.$$

By (1.12) and (4.1) we get that for any  $u \in Y_k$

$$\begin{aligned} \mathcal{J}_K(u) &\leq \frac{1}{2} \|u\|_{X_0}^2 - a_3 \|u\|_{L^\mu(\Omega)}^\mu + a_4 |\Omega| \\ &\leq \frac{1}{2} \|u\|_{X_0}^2 - \hat{C}_{k, \mu} \|u\|_{X_0}^\mu + a_4 |\Omega| \end{aligned} \tag{4.5}$$

for a suitable positive constant  $\hat{C}_{k, \mu}$  depending on  $k$  and  $\mu$ . As a consequence of (4.5), for any  $u \in Y_k$  with  $\|u\|_{X_0} = r_k$  we get that

$$\mathcal{J}_K(u) \leq 0,$$

provided  $r_k > 0$  is large enough, due to the fact that  $\mu > 2$ . Thus, step 1 is proved.

**Step 2.** There exists  $\gamma_k > 0$  such that

$$b_k = \inf \left\{ \mathcal{J}_K(u) : u \in Z_k, \|u\|_{X_0} = \gamma_k \right\} \rightarrow +\infty \quad \text{as } k \rightarrow +\infty.$$

By (1.8) and integrating, it is easy to see that (3.4) holds true, and so, as a consequence, we get that there exists a constant  $C > 0$  such that

$$|F(x, t)| \leq C(1 + |t|^q) \tag{4.6}$$

for any  $x \in \bar{\Omega}$  and  $t \in \mathbb{R}$ . Then, by (4.6) for any  $u \in Z_k \setminus \{0\}$ , we obtain

$$\begin{aligned} \mathcal{J}_K(u) &\geq \frac{1}{2} \|u\|_{X_0}^2 - C \|u\|_{L^q(\Omega)}^q - C |\Omega| \\ &= \frac{1}{2} \|u\|_{X_0}^2 - C \left\| \frac{u}{\|u\|_{X_0}} \right\|_{L^q(\Omega)}^q \|u\|_{X_0}^q - C |\Omega| \\ &\geq \frac{1}{2} \|u\|_{X_0}^2 - C \beta_k^q \|u\|_{X_0}^q - C |\Omega| \\ &= \|u\|_{X_0}^2 \left( \frac{1}{2} - C \beta_k^q \|u\|_{X_0}^{q-2} \right) - C |\Omega|, \end{aligned} \tag{4.7}$$

where  $\beta_k$  is defined as in lemma 6. Choosing

$$\gamma_k = (qC\beta_k^q)^{-1/(q-2)},$$

it is easy to see that  $\gamma_k \rightarrow +\infty$  as  $k \rightarrow +\infty$ , thanks to lemma 6 and the fact that  $q > 2$ . As a consequence of this and by (4.7) we get that for any  $u \in Z_k$  with  $\|u\|_{X_0} = \gamma_k$

$$\mathcal{J}_K(u) \geq \|u\|_{X_0}^2 \left( \frac{1}{2} - C \beta_k^q \|u\|_{X_0}^{q-2} \right) - C |\Omega| = \left( \frac{1}{2} - \frac{1}{q} \right) \gamma_k^2 - C |\Omega| \rightarrow +\infty$$

as  $k \rightarrow +\infty$ . Thus, step 2 is fulfilled.

The proof of theorem 1 under assumption (a) is complete.  $\square$

We notice that the Ambrosetti–Rabinowitz condition (1.9) was used just for proving step 1 in the verification of the geometric structure of the functional  $\mathcal{J}_K$  (actually we used (1.12)). While in the proof of step 2 the main tools were the assumption (1.8) and the Sobolev embedding theorems (see lemma 6).

4.2. Proof of theorem 1 under assumption (b)

By proposition 3 and (1.11), we have that  $\mathcal{J}_K$  satisfies the Cerami condition (and so the Palais–Smale condition) and  $\mathcal{J}_K(-u) = \mathcal{J}_K(u)$  for any  $u \in X_0$ . The verification of the geometric assumption (ii) of the Fountain Theorem follows as in step 2 in section 4.1. It remains to verify the condition (i). At this purpose we will use the finite-dimension of the linear subspace  $Y_k$  and assumption (1.13).

Indeed, for any  $k \in \mathbb{N}$ , by (1.13) there exists  $\delta_k > 0$  such that

$$F(x, t) \geq \frac{1}{C_k^2} |t|^2 \text{ for any } x \in \overline{\Omega} \text{ and any } t \in \mathbb{R} \text{ with } |t| > \delta_k, \tag{4.8}$$

where  $C_k := C_{k,2}$ , being  $C_{k,2}$  the positive constant given in (4.1) with  $q = 2$ . Moreover, by Weierstrass Theorem, we have that

$$F(x, t) \geq m_k := \min_{x \in \overline{\Omega}, |t| \leq \delta_k} F(x, t) \text{ for any } x \in \overline{\Omega} \text{ and any } t \in \mathbb{R} \text{ with } |t| \leq \delta_k. \tag{4.9}$$

Note that  $m_k \leq 0$ , since  $F(x, 0) = 0$  for any  $x \in \overline{\Omega}$ . By (4.8) and (4.9), it is easy to see that

$$F(x, t) \geq \frac{1}{C_k^2} |t|^2 - B_k \text{ for any } (x, t) \in \overline{\Omega} \times \mathbb{R}$$

for a suitable positive constant  $B_k$  (say,  $B_k \geq \delta_k^2/C_k^2 - m_k$ ).

As a consequence of this and by (4.1), for any  $u \in Y_k$  we have

$$\begin{aligned} \mathcal{J}_K(u) &= \frac{1}{2} \|u\|_{X_0}^2 - \int_{\Omega} F(x, u(x)) \, dx \\ &\leq \frac{1}{2} \|u\|_{X_0}^2 - \frac{1}{C_k^2} \|u\|_{L^2(\Omega)}^2 + B_k |\Omega| \\ &\leq \frac{1}{2} \|u\|_{X_0}^2 - \|u\|_{X_0}^2 + B_k |\Omega| \\ &= -\frac{1}{2} \|u\|_{X_0}^2 + B_k |\Omega|, \end{aligned} \tag{4.10}$$

so that, when  $\|u\|_{X_0} = r_k$  it follows that

$$\mathcal{J}_K(u) \leq 0,$$

provided  $r_k > 0$  is large enough. This proves that  $\mathcal{J}_K$  satisfies condition (i) of the Fountain Theorem and this ends the proof of theorem 1 under assumption (b).

4.3. Proof of theorem 1 under assumption (c)

The functional  $\mathcal{J}_K$  satisfies the Cerami condition by proposition 5, and so also the Palais–Smale assumption is verified. Moreover,  $\mathcal{J}_K(-u) = \mathcal{J}_K(u)$  for any  $u \in X_0$ , thanks to (1.11).

As for the geometric features of  $\mathcal{J}_K$ , condition (ii) of the Fountain Theorem follows as in step 2 of section 4.1. While condition (i) can be proved as in section 4.2. Hence, the proof of theorem 1 under assumption (c) is complete.



## Acknowledgments

The first author was supported by the Research Foundation of Heilongjiang Educational Committee (No. 12541667). The second and the third authors were supported by the INdAM-GNAMPA Project 2014 *Proprietà geometriche e analitiche per problemi non-locali*. The third author was supported by the MIUR National Research Project *Variational and Topological Methods in the Study of Nonlinear Phenomena* and by the ERC grant  $\epsilon$  (*Elliptic Pde's and Symmetry of Interfaces and Layers for Odd Nonlinearities*). We would like to thank the referee for her/his useful and interesting comments and suggestions.

## References

- [1] Adams R A 1975 *Sobolev Spaces* (New York: Academic)
- [2] Alves C O and Liu S B 2010 On superlinear  $p(x)$ -Laplacian equations in  $\mathbb{R}^N$  *Nonlinear Anal.* **73** 2566–79
- [3] Ambrosetti A and Rabinowitz P 1973 Dual variational methods in critical point theory and applications *J. Funct. Anal.* **14** 349–81
- [4] Autuori G and Pucci P 2013 Elliptic problems involving the fractional Laplacian in  $\mathbb{R}^N$  *J. Differ. Equ.* **25** 2340–62
- [5] Barrios B, Colorado E, Servadei R and Soria F A critical fractional equation with concave–convex power nonlinearities to appear in *Ann. Inst. H. Poincaré Anal. Non Linéaire*
- [6] Bartsch T 1993 Infinitely many solutions of a symmetric Dirichlet problem *Nonlinear Anal.* **20** 1205–16
- [7] Brezis H 1983 *Analyse Fonctionnelle. Théorie et Applications* (Paris: Masson)
- [8] Cabré X and Tan J 2010 Positive solutions of nonlinear problems involving the square root of the Laplacian *Adv. Math.* **224** 2052–93
- [9] Caffarelli L and Silvestre L 2007 An extension problem related to the fractional Laplacian *Commun. Partial Differ. Equ.* **32** 1245–60
- [10] Caffarelli L and Silvestre L 2009 Regularity theory for fully nonlinear integro-differential equations *Commun. Pure Appl. Math.* **62** 597–638
- [11] Cerami G 1978 An existence criterion for the critical points on unbounded manifolds *Ist. Lomb. Accad. Sci. Lett. Rend. A* **112** 332–6
- [12] Cerami G 1980 On the existence of eigenvalues for a nonlinear boundary value problem *Ann. Mat. Pura Appl.* **124** 161–79
- [13] Costa D G and Magalh C A 1994 Variational elliptic problems which are nonquadratic at infinity *Nonlinear Anal.* **23** 1401–12
- [14] Di Nezza E, Palatucci G and Valdinoci E 2012 Hitchhiker's guide to the fractional Sobolev spaces *Bull. Sci. Math.* **136** 521–73
- [15] Fang F and Liu S B 2009 Nontrivial solutions of superlinear  $p$ -Laplacian equations *J. Math. Anal. Appl.* **351** 138–46
- [16] Ferrara M, Molica Bisci G and Zhang B 2014 Existence of weak solutions for non-local fractional problems via Morse theory *Discrete Contin. Dyn. Syst. B* **19** 2493–9
- [17] Fiscella A, Servadei R and Valdinoci E 2013 A resonance problem for non-local elliptic operators *Z. Anal. Anwendungen* **32** 411–31
- [18] Fiscella A and Valdinoci E 2014 A critical Kirchhoff type problem involving a nonlocal operator *Nonlinear Anal.* **94** 156–70
- [19] Jeanjean L 1999 On the existence of bounded Palais-Smale sequences and application to a Landesman–Lazer type problem set on  $\mathbb{R}^N$  *Proc. R. Soc. Edinburgh A* **129** 787–809
- [20] Liu S B 2010 On ground states of superlinear  $p$ -Laplacian equations in  $\mathbb{R}^N$  *J. Math. Anal. Appl.* **361** 48–58
- [21] Liu S B 2010 On superlinear problems without Ambrosetti–Rabinowitz condition *Nonlinear Anal.* **73** 788–95
- [22] Liu S B and Liu S J 2003 Infinitely many solutions for a superlinear elliptic equation *Acta. Math. Sin.* **46** 625–30
- [23] Miyagaki O and Souto M 2008 Superlinear problems without Ambrosetti and Rabinowitz growth condition *J. Differ. Equ.* **245** 3628–38
- [24] Molica Bisci G 2014 Fractional equations with bounded primitive *Appl. Math. Lett.* **27** 53–58
- [25] Molica Bisci G 2014 Sequences of weak solutions for fractional equations *Math. Res. Lett.* **21** 1–13
- [26] Molica Bisci G and Pansera B A 2014 Three weak solutions for nonlocal fractional equations *Adv. Nonlinear Stud.* **14** 591–601
- [27] Pucci P and Saldi S Critical stationary Kirchhoff equations in  $\mathbb{R}^N$  involving nonlocal operators to appear in *Rev. Mat. Iberoam*

AQ1

AQ2

- [28] Rabinowitz P H 1986 *Minimax Methods in Critical Point Theory with Applications to Differential Equations* (CBMS Regional Conf. Series in Mathematics vol 65) (Providence, RI: American Mathematical Society)
- [29] Ros-Oton X and Serra J 2014 The Dirichlet problem for the fractional Laplacian: regularity up to the boundary *J. Math. Pures Appl.* **101** 275–302
- [30] Schechter M and Zou W 2004 Superlinear problems *Pac. J. Math.* **214** 145–60
- [31] Servadei R 2013 The Yamabe equation in a non-local setting *Adv. Nonlinear Anal.* **2** 235–70
- [32] Servadei R and Valdinoci E 2013 Lewy–Stampacchia type estimates for variational inequalities driven by (non)local operators *Rev. Mat. Iberoam.* **29** 1091–126
- [33] Servadei R and Valdinoci E 2012 Mountain pass solutions for non-local elliptic operators *J. Math. Anal. Appl.* **389** 887–98
- [34] Servadei R and Valdinoci E 2013 Variational methods for non-local operators of elliptic type *Discrete Contin. Dyn. Syst.* **33** 2105–37
- [35] Struwe M 1990 *Variational Methods: Applications to Nonlinear Partial Differential Equations and Hamiltonian Systems* (Ergebnisse der Mathematik und ihrer Grenzgebiete vol 3) (Berlin: Springer)
- [36] Tan J 2011 The Brezis–Nirenberg type problem involving the square root of the Laplacian *Calc. Var. Partial Differ. Equ.* **36** 21–41
- [37] Willem M 1996 *Minimax Theorems* (Boston: Birkhäuser)
- [38] Willem M and Zou W 2003 On a Schrödinger equation with periodic potential and spectrum point zero *Indiana Univ. Math. J.* **52** 109–32
- [39] Zou W M 2001 Variant fountain theorems and Their applications *Manuscr. Math.* **104** 343–58

## QUERIES

### Page 17

AQ1

Please check the details for any journal references that do not have a blue link as they may contain some incorrect information. Pale purple links are used for references to arXiv e-prints.

### Page 17

AQ2

Please update the year and publication details if appropriate in references [[5](#),[27](#)].